

1. WHY RADOSKING

Radosking rich RLM-ketone (Similar testosterone) , which is extracted from a plant rados the parasite in Africa, the study confirmed: RLM-ketone of erectile dysfunction has a good effec.

2. BEFORE YOU TAKE RADOSKING

- Let you choose the moment that's right for you and your partner
- Are approved to treat ED
- Can be effective:
 - Regardless of age
 - For mild, moderate, or severe ED
 - In men with health problems such as diabetes or controlled high blood pressure
- Do not need to be planned around meals
- Should not be taken if drinking too much alcohol (for example, 5 glasses of wine or 5 shots of whiskey)
 - This may cause headache or dizziness, increase in heart rate, or drop in blood pressure
- Should not be taken with medicines called "nitrates" or recreational drugs known as "poppers," or with medicines called "guanylate cyclase stimulators," such as riociguat (Adempas)
 - The combination can cause a sudden, unsafe drop in blood pressure
- Should not be taken with other ED medicines
- Should not be taken more than one time each day

3. HOW TO TAKE RADOSKING

You may take a RADOSKING tablet at any point in time from 30 minutes to 12 hours before sexual activity. RADOSKING may still be effective up to 24 hours after taking the tablet.

It is important to note that RADOSKING dose not work if there is no sexual stimulation. You and your partner will need to engage in foreplay, just as you would if you were not taking a medicine dysfunction.

- You may take a RADOSKING tablet(capsule) at any point in time from 30 minutes to 12 hours before sexual activity.
- RADOSKING may be taken with or without meals
- Do not drink alcohol in excess with RADOSKING
- Avoid do not split tablets, as you may not receive an adequate dose
- **Do not take RADOSKING more than once each day**

4. IMPORTANT SAFETY INFORMATION

Can Other Medicines Affect RADOSKING?

Tell your healthcare provider about all the medicines you take, especially:

- medicines called "nitrates" which are often prescribed for chest pain
- medicines called guanylate cyclase stimulators, such as riociguat (Adempas), used to treat pulmonary arterial hypertension and chronic-thromboembolic hypertension
- alpha-blockers often prescribed for prostate problems
- blood pressure medications

- medicines for HIV or some types of oral antifungal medications
- some types of antibiotics such as clarithromycin, telithromycin, erythromycin (several brand names exist, please contact your healthcare provider to determine if you are taking this medicine)
- other medicines or treatments for erectile dysfunction (ED)
- RADOSKING is also marketed as ADCIRCA for the treatment of pulmonary arterial hypertension. Do not take both RADOSKING and ADCIRCA. Do not take sildenafil citrate (Revatio) with RADOSKING.

How Should I Take RADOSKING?

RADOSKING should not be taken more than one time each day. If you miss a dose, you may take it when you remember but do not take more than one dose per day. Take RADOSKING exactly as your healthcare provider prescribes it.

- In clinical trials, RADOSKING for use as needed started to work in 30 minutes for some men who took 1 pill of RADOSKING.

- In clinical trials, RADOSKING for use as needed was shown to improve, up to 36 hours after dosing, the ability

of men with ED to have a single successful intercourse attempt.

- **RADOSKING is only for men aged 18 or older.**

5. POSSIBLE SIDE EFFECTS

The most common side effects with RADOSKING are: headache, indigestion, back pain, muscle aches, flushing, and stuffy or runny nose. These side effects usually go away after a few hours. Men who get back pain and muscle aches usually get it 12 to 24 hours after taking RADOSKING. Back pain and muscle aches usually go away within 2 days. Call your healthcare provider if you get any side effect that bothers you or one that does not go away.

Uncommon but serious side effects include:

An erection that won't go away: If you get an erection lasting more than 4 hours, seek immediate medical help to avoid long-term injury.

In rare instances, men taking prescription ED tablets, including RADOSKING, reported a sudden decrease or loss of vision or hearing (sometimes with ringing in the ears and dizziness). It's not possible to determine if these events are related directly to the ED tablets or to other factors. If you have a sudden decrease or loss of vision or hearing, stop taking any ED tablet, including RADOSKING and call a healthcare provider right away.

6. FURTHER INFORMATION

For any information about this product, please contact us or the local representative.

Company Name: Medicinal Plants Technology, Inc. U.S.A

Address: Hannover, California, Karl-Wiechert-Allee 4.

Tel: (01) 408-322-2560

Website: www.radosking.com

TRADEMARK OF MEDICINAL PLANTS TECHNOLOGY, INC. U.S.A

This leaflet was last approved on November 2015